19

.....

ACIDS, BASES, AND SALTS

SECTION 19.1 ACID-BASE THEORIES (pages 587–593)

This section compares and contrasts acids and bases as defined by the theories of Arrhenius, Brønsted-Lowry, and Lewis. It also identifies conjugate acid–base pairs in acid–base reactions.

Properties of Acids and Bases (pages 587–588)

- 1. Circle the letters of all the terms that complete the sentence correctly. The properties of acids include ______.
 - a. reacting with metals to produce oxygen
 - **b.** giving foods a sour taste
 - c. forming solutions that conduct electricity
 - d. causing indicators to change color
- 2. Bases are compounds that react with acids to form _____

and a(n) _____

- **3.** Circle the letters of all the terms that complete the sentence correctly. The properties of bases include ______.
 - a. tasting bitter
 - **b.** feeling slippery
 - c. changing the color of an indicator
 - d. always acting as a strong electrolyte

bonds are ionizable.

Arrhenius Acids and Bases (pages 588–590)

4. Match the number of ionizable hydrogens with the type of acid.

 _____ one
 a. diprotic

 _____ two
 b. triprotic

- _____ three c. monoprotic
- 5. Is the following sentence true or false? Only the hydrogens in weak polar
- 6. Hydrogen is joined to a very ______ element in a very polar bond.
- 7. Alkali metals react with water to produce ______ solutions.

8. How do concentrated basic solutions differ from other basic solutions?

	Is the following sentence true or false? Some of the acids and bases included in
	the Arrhenius theory are not acids and bases according to the Brønsted-Lowry
	theory
•	Is the following sentence true or false? A conjugate acid is the particle formed
	when a base gains a hydrogen ion
•	A conjugate is the particle that remains when an acid has donated a hydrogen ion.
•	What is a conjugate acid–base pair?
•	A substance that can act as both an acid and a base is said to be
	In a reaction with HCL is water an acid or a base?

17. A Lewis base is a substance that can ______ a pair of electrons to form a covalent bond.

- **18.** Is the following sentence true or false? All the acids and bases included in the Brønsted-Lowry theory are also acids and bases according to the Lewis theory.
- **19.** Complete this table of acid-base definitions.

Acid–Base Definitions		
Туре	Acid	Base
Brønsted-Lowry		H⁺ acceptor
	electron-pair acceptor	
	H ⁺ producer	

SECTION 19.2 HYDROGEN IONS AND ACIDITY (pages 594–604)

This section classifies solutions as neutral, acidic, or basic, given the hydrogenion or hydroxide-ion concentration. It explains how to convert hydrogen-ion concentrations into pH values and hydroxide-ion concentrations into pOH values.

Hydrogen lons from Water (pages 594-595)

- 1. What does a water molecule that loses a hydrogen ion become?
- 2. What does a water molecule that gains a hydrogen ion become?

3. The reaction in which water molecules produce ions is called the

_____ of water.

4. In water or aqueous solution, ______ are always joined to

_____ as hydronium ions (H_3O^+).

5. Is the following sentence true or false? Any aqueous solution in which [H⁺] and [OH⁻] are equal is described as a neutral solution.

Ion Product Constant for Water (pages 595–596)

6. What is the ion-product constant for water (*K*_w)? Give the definition, the expression, and the value.

7. A(n) solution is one in which [H ⁺] is greater than [OI

A(n) ______ solution is one in which $[H^+]$ is less than $[OH^-]$.

8. Match the type of solution with its hydrogen-ion concentration.

acidic	a. less than 1.0 $ imes$ 10 ⁻⁷ M
neutral	b. greater than $1.0 \times 10^{-7} M$
basic	c. $1.0 imes10^{-7}M$

The pH Concept (pages 596–600)

9. The _____ of a solution is the negative logarithm of the hydrogen-ion concentration.

10. Match the type of solution with its pH.

acidic	a. pH > 7.0
neutral	b. pH = 7.0
basic	c. pH < 7.0

- 11. Look at Table 19.5 on page 598. What is the approximate [H⁺], the [OH⁻], and the pH of washing soda?
- **12.** The pOH of a solution is the negative logarithm of the

_____ concentration.

- **13.** What is the pOH of a neutral solution?
- **14.** For pH calculations, in what form should you express the hydrogen-ion concentration?
- **15.** Look at the pH scale below. Label where you would find acids, bases, and neutral solutions.



16. Is the following sentence true or false? Most pH values are whole numbers.

17. If [H⁺] is written in scientific notation but its coefficient is not 1, what do you need to calculate pH?

18. Is the following sentence true or false? You can calculate the hydrogen-ion concentration of a solution if you know the pH. _____

Measuring pH (pages 600–603)

19. When do you use indicators and when do you use a pH meter to measure pH?

20. Why is an indicator a valuable tool for measuring pH?

- **21.** Why do you need many different indicators to span the entire pH spectrum?
- **22.** Look at the figure below. Fill in the missing pH color change ranges for the indicators.



23. List three characteristics that limit the usefulness of indicators.



- 24. How accurate are measurements of pH obtained with a pH meter?
- 25. What is the pH of each of the following solutions?
 - **a.** water _____
 - **b.** vinegar _____
 - c. milk of magnesia _____
- **26.** Is the following sentence true or false? Measurements of pH obtained with a pH meter are typically accurate to within 0.001 pH unit of the true pH.

SECTION 19.3 STRENGTHS OF ACIDS AND BASES (pages 605–611)

This section defines strong acids and weak acids, and then explains how to calculate an acid dissociation constant. It describes how acids and bases are arranged by strength according to their dissociation constants (K_a) and (K_b).

Strong and Weak Acids and Bases (pages 605–609)

1. What factor is used to classify acids as strong or weak?

2.	Strong acids are ionized in aqueous solution; weak
	acids ionize in aqueous solution.
3.	Look at Table 19.6 on page 605. Which acid is the weakest acid in the table? Which base is the weakest base?
4.	What do you use to write the equilibrium-constant expression?
5.	An acid dissociation constant (K_a) is the ratio of the concentration of the
	form of an acid to the concentration of the
	form.
6.	What is another name for dissociation constants?
7.	Is the following sentence true or false? The stronger an acid is, the smaller its
	<i>K</i> _a value
8.	A diprotic acid has dissociation constants.

- **9.** Look at Table 19.7 on page 607. What is the second dissociation constant for the triprotic phosphoric acid? ______
- **10.** Weak bases react with water to form the hydroxide ion and the ______ of the base.
- **11.** A base dissociation constant (K_b) is the ratio of the concentration of the

_____ times the concentration of the hydroxide

ion to the concentration of the ______.

- **12.** What does the magnitude of the base dissociation constant (K_b) indicate?
- **13.** The words *concentrated* and *dilute* indicate how much acid or base is in solution.
- **14.** Is the following sentence true or false? The words strong or weak refer to the extent of ionization or dissociation of an acid or base.

Calculating Dissociation Constants (pages 609–610)

- **15.** Is the following sentence true or false? You can calculate the acid dissociation constant (*K*_a) of a weak acid from experimental data.
- **16.** To measure the equilibrium concentrations of all substances present at equilibrium for a weak acid, what two conditions must you know?



Reading Skill Practice

By looking carefully at photographs and drawings in textbooks, you can better understand what you have read. Look carefully at Figure 19.16 on page 606. What important idea does this drawing communicate? Do your work on a separate sheet of paper.

SECTION 19.4 NEUTRALIZATION REACTIONS (pages 612–616)

This section explains how acid–base titration is used to calculate the concentration of an acid or a base. It also explains the concept of equivalence in neutralization reactions.

Acid-Base Reactions (pages 612-613)

1.	Is the following sentence true or false? Acids react with compounds containing
	hydroxide ions to form water and a salt.

2. What does the reaction of an acid with a base produce?

3. In general, reactions in which an acid and a base react in an aqueous solution

to produce a salt and water are called ______ reactions.

- **4.** Look at Table 19.9 on page 613. Circle the letter of the salt that is used for photographic emulsions.
 - **a.** calcium chloride **c.** silver bromide
 - **b.** potassium chloride **d.** sodium chloride
- 5. Salts are compounds consisting of a(n) ______ from an acid
 - and a(n) ______ from a base.

Titration (pages 613–616)

- 6. How can you determine the concentration of an acid or base in a solution?
- 7. Complete the flow chart below showing the steps of a neutralization reaction.

A measured volume of an acid solution of concentration is added to a flask.	
Several drops of the while the flask is gently swirled.	_ are added to the solution
¥	
Measured volumes of a base of mixed into the acid until the indicator cha	nges

8. The process of adding a known amount of solution of known concentration to

determine the concentration of another solution is called ______.

9. What is the solution of known concentration called?

SECTION 19.5 SALTS IN SOLUTION (pages 618–622)

This section demonstrates with equations how buffers resist changes in pH. It also explains how to calculate the solubility product constant of a slightly soluble salt.

Salt Hydrolysis (pages 618–620)

- 1. What is salt hydrolysis? _____
- **2.** Complete this table of the rules for hydrolysis of a salt.

Reactants	Products
acid + base	Neutral solution
Strong acid + Weak base	solution
acid + base	Basic solution

Buffers (pages 620-622)

- 3. What are buffers? _____
- 4. A buffer is a solution of a ______ acid and one of its salts,

or a solution of a _____ base and one of its salts.

 Is the following sentence true or false? The buffer capacity is the amount of acid or base that can be added to a buffer solution before a significant change in pH occurs.

GUIDED PRACTICE PROBLEMS

EXTRA PRACTICE PROBLEM (similar to Practice Problem 13, page 600)

13. Find the value of $[OH^{-}]$ for a solution with a pH of 8.00.

GUIDED PRACTICE PROBLEM 16b (page 601)

16b. Calculate the pH of this solution: $[H^+] = 8.3 \times 10^{-10} M$.

Step 1. Identify the known and unknown values.	Known [H ⁺] = $\longrightarrow 10^{-10} M$	Unknown pH = ?
Step 2. Substitute values into the pH equation.	$pH = -\log [H^+]$ $= -\log (8.3 \times \square)$)
Step 3. The logarithm of a product equals the sum of the logs of its factors.	= - (log + log)
Step 4. Evaluate $\log 8.3$ by using a calculator. Evaluate $\log 10^{-10}$ by using the definition of logarithm.	= -(0.919 +)	
Step 5. Add and simplify. Write your answer with two significant figures to the right of the decimal point.	= -(-9.081) =	

GUIDED PRACTICE PROBLEM 22 (page 610)

22. For a solution of methanoic acid exactly 0.1 *M*, $[H^+] = 4.2 \times 10^{-3} M$. Calculate the K_a of methanoic acid.

 $K_{\rm a} =$

Analyze

Step 1. What is known about the acid?

Step 2. What is the unknown? _____

Step 3. What is the expression for the K_a of methanoic acid?

Solve

Step 4. What expression can you write to find the equilibrium concentration of HCOOH?

Step 5. Substitute values into the formula for K_a and solve.

Analyze

Step 6. Look at Table 19.7 on page 607. Explain why your answer is reasonable.