

Name: \_\_\_\_\_

Date: \_\_\_\_\_ Period: \_\_\_\_\_

1<sup>st</sup> & 2<sup>nd</sup> Derivative Test Review Sheet

Calculus Honors

**Review Sheet: Applications of Derivatives**

In the examples below, locate the absolute extrema of a function on the indicated closed interval.

1.  $f(x) = \frac{2x+5}{3}, [0,5]$

2.  $f(x) = -x^2 + 3x, [0,3]$



3.  $y = 3x^{2/3} - \frac{3}{2}x^2, [-1,2]$

4.  $h(s) = \frac{1}{s-2}, [0,1]$

5.  $f(x) = 2 \sin x - \cos 2x, [0,2\pi]$

6.  $g(x) = \sec x, \left[-\frac{\pi}{6}, \frac{\pi}{3}\right]$

In the examples below, decide whether each labeled point is an absolute maximum or minimum, a relative maximum or minimum, or neither.



In the examples below, determine whether Rolle's Theorem can be applied to  $f$  on the closed interval  $[a,b]$ . If Rolle's Theorem can be applied, find all values  $c$  in the open interval  $(a, b)$  such that  $f'(c) = 0$ .

9.  $f(x) = x^2 - 2x, [0,2]$

10.  $f(x) = \sin x, [0,2\pi]$

11.  $f(x) = \frac{x^2-1}{x}, [-1,1]$

In the examples below, use the Mean Value Theorem to solve the word problems.

12. When an object is removed from a furnace and placed in an environment with a constant temperature of  $90^\circ\text{F}$ , its core temperature is  $1500^\circ\text{F}$ . Five hours later the core temperature is  $390^\circ\text{F}$ . Explain why there must exist a time in the interval when the temperature is decreasing at a rate of  $222^\circ\text{F}$  per hour.

13. At 9:13 a sports car is traveling 35 miles per hour. Two minutes later, the car is traveling at 85 miles per hour. Prove that at some time during this two-minute interval, the car's acceleration is exactly 1500 miles per hour squared.

In the examples below, (a) find the critical numbers of  $f$  (if any), (b) find the open interval(s) on which the function is increasing or decreasing, and (c) apply the First Derivative Test to identify all relative extrema.

14.  $f(x) = x^2 - 6x$

15.  $f(x) = 2x^3 + 3x^2 - 12x$

16.  $f(x) = \frac{x^5 - 5x}{5}$

In the examples below, consider the function on the interval  $(0, 2\pi)$ , (a) find the critical numbers of  $f$  (if any), (b) find the open interval(s) on which the function is increasing or decreasing, and (c) apply the First Derivative Test to identify all relative extrema.

17.  $f(x) = \sin x + \cos x$

18.  $f(x) = \sqrt{3} \sin x + \cos x$

19.  $f(x) = \sin^2 x + \sin x$

In the examples below, find the points of inflection and discuss the concavity of the graph of the function.

20.  $f(x) = x^3 - 6x^2 + 12x$

21.  $f(x) = 2x^3 - 3x^2 - 12x + 5$

22.  $f(x) = \sin \frac{x}{2}, [0, 4\pi]$

In the examples below, find all relative extrema. Use the Second Derivative Test.

23.  $f(x) = x^4 - 4x^3 + 2$

24.  $f(x) = \sqrt{x^2 + 1}$

25.  $f(x) = \cos x - x, [0, 4\pi]$

**Answer Key:**

- 1.) (0, 5/3) minimum; (5,5) maximum                      2.) (0,0)& (3, 0) minimum; (3/2, 9/4) maximum
- 3.) (2, -1.2378) minimum; (-1, 1.5) maximum                      4.) (0, -1/2) max; (1,-1) min
- 5.) (0,-1),  $(\frac{3\pi}{2}, -1)$ ,  $(2\pi, -1)$  min;  $(\frac{\pi}{2}, 3)$  max                      6.) (0,1) min;  $(\frac{\pi}{3}, 2)$  max
- 7.) A: relative max; B: relative min; C: absolute max; D: absolute min
- 8.) A: absolute max; B: neither; C: relative max; D: relative min
- 9.) Rolle's Theorem can be applied;  $c = 1$                       10.) Rolle's Theorem can be applied;  $c = \{\frac{\pi}{2}, \frac{3\pi}{2}\}$
- 11.) Rolle's Theorem cannot be applied because it's not a continuous function (discontinuous @  $x = 0$ )
- 12.) When the Mean Value Theorem is applied, the answer is  $-222^{\circ}\text{F}/\text{hour}$ . Since that is the average throughout the interval, there has to be a least one time when it's decreasing at that rate.
- 13.) When the Mean Value Theorem is applied, the answer is  $1500 \text{ miles per hour}^2$ . Since that is the average throughout the interval, there has to be a least one time when it's increasing at that rate.
- 14.) decreasing:  $(-\infty, 3)$ ; increasing:  $(3, \infty)$ ; minimum @  $(3, -9)$
- 15.) increasing:  $(-\infty, -2)$  &  $(1, \infty)$ ; decreasing:  $(-2, 1)$ ; max @  $(-2, 20)$ ; min @  $(1, -7)$
- 16.) increasing:  $(-\infty, -1)$  &  $(1, \infty)$ ; decreasing:  $(-1, 1)$ ; max @  $(-1, 0.8)$ ; min @  $(1, -0.8)$
- 17.) increasing:  $(0, \frac{\pi}{4})$  &  $(\frac{5\pi}{4}, 2\pi)$ ; decreasing:  $(\frac{\pi}{4}, \frac{5\pi}{4})$ ; max @  $(\frac{\pi}{4}, \sqrt{2})$ , min @  $(\frac{5\pi}{4}, -\sqrt{2})$
- 18.) increasing:  $(0, \frac{\pi}{3})$  &  $(\frac{4\pi}{3}, 2\pi)$ ; decreasing:  $(\frac{\pi}{3}, \frac{4\pi}{3})$ ; max @  $(\frac{\pi}{3}, 2)$ , min @  $(\frac{4\pi}{3}, -2)$
- 19.) increasing:  $(0, \frac{\pi}{2})$ ,  $(\frac{7\pi}{6}, \frac{3\pi}{2})$ ,  $(\frac{11\pi}{6}, 2\pi)$ ; decreasing:  $(\frac{\pi}{2}, \frac{7\pi}{6})$ ,  $(\frac{3\pi}{2}, \frac{11\pi}{6})$ ; max:  $(\frac{\pi}{2}, 2)$ ,  $(\frac{3\pi}{2}, 0)$ ;  
min:  $(\frac{7\pi}{6}, -\frac{1}{4})$ ,  $(\frac{11\pi}{6}, -\frac{1}{4})$